

Total No. of Printed Pages—7

3 SEM TDC MTMH (CBCS) C 5

2 0 2 1

(Held in January/February, 2022)

MATHEMATICS

(Core)

Paper : C-5

(Theory of Real Functions)

Full Marks : 80

Pass Marks : 32

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

- ~~1.~~ (a) Define limit of function at a point. 1
- ~~(b)~~ Evaluate the following limits (any one) : 2

(i) $\lim_{x \rightarrow 2} \sqrt{\frac{2x+1}{x+3}}$

(ii) $\lim_{x \rightarrow 1} \frac{x-1}{\sqrt{x+3}-2}$

(2)

- (c) If $f : A \rightarrow R$ and if c is a cluster point of A , then prove that f can have only one limit at c . 3

2. (a) Write the type of discontinuity if

$$\lim_{x \rightarrow c^+} f(x) \neq \lim_{x \rightarrow c^-} f(x)$$
 1

- (b) When does a function f continuous on a set? 2

- (c) Investigate for the point of discontinuity : 4

$$f(x) = \begin{cases} 1 ; & \text{if } x \text{ is rational} \\ 0 ; & \text{if } x \text{ is irrational} \end{cases}$$

Or

Let $A, B \subseteq R$ and let $f : A \rightarrow R$ and $g : B \rightarrow R$ be functions such that $f(A) \subseteq B$. If f is continuous at a point $c \in A$ and g is continuous at $b = f(c) \in B$; then prove that composition $g \circ f : A \rightarrow R$ is continuous at c .

(3)

- (d) Let $A \subseteq \mathbb{R}$, let $f : A \rightarrow \mathbb{R}$ and let $|f|$ be defined by $|f|(x) = |f(x)|$ for $x \in A$ and f is continuous at a point $c \in A$. Prove that $|f|$ is continuous at c . 3

Or

Discuss the continuity of $f(x) = |x - 1| + |x - 2|$ in the interval $[0, 3]$.

3. (a) State location of roots theorem. 1
- (b) State and prove intermediate value theorem. 4
- (c) Find the roots of the equation $x^3 - x - 1 = 0$ between 1 and 2 by using location of roots (bisection method) theorem. 4

Or

Let I be a closed bounded interval and let $f : I \rightarrow \mathbb{R}$ be continuous on I , then prove that the set $f(I) = \{f(x) : x \in I\}$ is a closed bounded interval.

4. (a) Write the non-uniformity continuity criteria (any one). 1

- (b) Show that a function $f : R \rightarrow R$ given by $f(x) = x^2$ is not uniformly continuous on R .

4

Or

If f and g are each uniformly continuous on R , then prove that composite function $f \circ g$ is also uniformly continuous on R .

5. (a) Find :

$$\frac{d}{dx}(\tan x^2)$$

1

- (b) State Caratheodory's theorem.

2

- (c) If f is continuous on the closed interval $I = [a, b]$ and f is differentiable on the open interval (a, b) and $f'(x) = 0$ for all $x \in (a, b)$, prove that f is constant on I .

3

6. (a) Define relative maximum and relative minimum at a point on an interval.

2

- (b) State and prove Rolle's theorem. 1+3=4

(5)

(c) Apply the mean value theorem to prove the following (any one) :

4

(i) $e^x \geq 1+x$ for $x \in R$

(ii) $\frac{b-a}{1+b^2} < \tan^{-1} b - \tan^{-1} a < \frac{b-a}{1+a^2}$
for $a < b$

7. (a) Show that $f(x) = x^3 - 3x^2 + 3x + 2$ is strictly increasing for every value of $x \in R$ except 1.

2

(b) Let $I \subseteq R$ be an interval, let $f : I \rightarrow R$, let $c \in I$ and assume that f has a derivative at c and $f'(c) > 0$, then there is a number $\delta > 0$. Prove that $f(x) > f(c)$ for $x \in I$ and $c < x < c + \delta$.

3

(c) Examine the validity of mean value theorem for the function $f(x) = 2x^2 - 7x + 10$ on $[2, 5]$.

4

Or

If f is differentiable on $I = [a, b]$ and if k is a number between $f'(a)$ and $f'(b)$, then prove that there exists at least one point c in (a, b) , where $f'(c) = k$.

8. (a) Write the remainder after n terms of Taylor's theorem in Lagrange's form. 1

(b) Write the statement of Cauchy's mean value theorem. 2

(c) Deduce from Cauchy's mean value theorem $f(b) - f(a) = \xi f'(\xi) \log \frac{b}{a}$, where $f(x)$ is continuous and differentiable in $[a, b]$ and $a < \xi < b$. 3

(d) State and prove Taylor's theorem with Cauchy's form of remainder. 6

Or

Find the approximate value of $\sqrt[3]{1+x}$, $x > -1$ by using Taylor's theorem with $n = 2$.

9. (a) Write the necessary condition for a function $f(x)$ to have relative extremum at $x = c$. 1

(b) Determine whether or not $x = 0$ is a point of relative extremum of $f(x) = \sin x - x$. 2

(c) Define convex function. 2

(7)

(d) Using Maclaurin's series, expand the following in an infinite series in powers of x (any two) : 4×2=8

(i) $\log(1+x)$

(ii) $\cos x$

(iii) $\frac{1}{ax+b}$
